Oxytocin enhances brain reward system responses in men viewing the face of their female partner

Dirk Scheele^{a,b,1}, Andrea Wille^{a,b,1}, Keith M. Kendrick^{c,1}, Birgit Stoffel-Wagner^d, Benjamin Becker^{a,b}, Onur Güntürkün^e, Wolfgang Maier^{a,f}, and René Hurlemann^{a,b,2}

Departments of ^aPsychiatry, ^bMedical Psychology, and ^dClinical Chemistry and Clinical Pharmacology, University of Bonn, 53105 Bonn, Germany; ^cKey Laboratory for Neuroinformation, School of Life Science and Technology, University of Electronic Science and Technology of China, 610054 Chengdu, China; ^eDepartment of Biopsychology, Ruhr University, 44780 Bochum, Germany; and ^fGerman Center for Neurodegenerative Diseases, 53175 Bonn, Germany

Edited by Angela Sirigu, Centre National de la Recherche Scientifique, Bron, France, and accepted by the Editorial Board November 6, 2013 (received for review July 26, 2013)

The biological mechanisms underlying long-term partner bonds in humans are unclear. The evolutionarily conserved neuropeptide oxytocin (OXT) is associated with the formation of partner bonds in some species via interactions with brain dopamine reward systems. However, whether it plays a similar role in humans has as yet not been established. Here, we report the results of a discovery and a replication study, each involving a double-blind, placebocontrolled, within-subject, pharmaco-functional MRI experiment with 20 heterosexual pair-bonded male volunteers. In both experiments, intranasal OXT treatment (24 IU) made subjects perceive their female partner's face as more attractive compared with unfamiliar women but had no effect on the attractiveness of other familiar women. This enhanced positive partner bias was paralleled by an increased response to partner stimuli compared with unfamiliar women in brain reward regions including the ventral tegmental area and the nucleus accumbens (NAcc). In the left NAcc, OXT even augmented the neural response to the partner compared with a familiar woman, indicating that this finding is partner-bond specific rather than due to familiarity. Taken together, our results suggest that OXT could contribute to romantic bonds in men by enhancing their partner's attractiveness and reward value compared with other women.

emotion | functional imaging | love | monogamy

Love and enduring romantic bonds can bring the elation of profound joy and pleasure but also, when broken, the deepest sorrow and despair. Although love is the source of a large variety of emotions and feelings and celebrated in all human cultures by countless works of art, as yet surprisingly little is known about the neurobiological underpinnings of long-term pair bonds (i.e., enduring attachments between sexual partners) in humans.

Indeed, very few other mammalian species $[\sim 3-5\% (1)]$ exhibit pair bonds, with the most investigated species being the prairie vole (Microtus ochrogaster) (2). In prairie voles, partner bonds are formed after the neuropeptides oxytocin (OXT) or arginine vasopressin (AVP) are released in the brain during mating. Their effects on bonding are mediated via interactions with dopamine (DA) release, particularly in the ventral tegmental area (VTA) and nucleus accumbens (NAcc). Furthermore, in different species of voles, a monogamous, as opposed to polygamous, pattern of behavior is associated with a higher density of dopamine D2 receptors (D2R) in the medial prefrontal cortex (mPFC) (3) and of OXT receptors (OXTR) in mPFC, NAcc, and caudate putamen (4, 5). In prairie voles, D2R activation in the NAcc facilitates a partner preference and bond formation in the absence of mating in both sexes (6), whereas the direct injection of OXT into the NAcc (7) does so in females and of AVP into the lateral septum of males (8). Conversely, the infusion of an AVP antagonist into the ventral pallidum of male prairie voles prevents partner preference formation (9). However, nonspecific intracerebroventricular administration of OXT

can induce a partner preference and bonding in both males and females (10).

These findings in voles are paralleled in humans by functional MRI (fMRI) studies showing that viewing the face of a romantic partner while recalling experiences with them activates rewardassociated regions such as the VTA, NAcc, caudate, and lentiform nuclei (11-14). Genetic variability in OXT influences DA signaling in brain reward centers (15), and there are also D2R-OXTR heteromers with facilitatory receptor-receptor interactions in the striatum (16). Moreover, numerous studies have reported prosocial effects of OXT including increased social affiliation and attachment (17-19). Plasma OXT levels can be elevated by several attachment-related experiences, including partner hugs (20), social support (21), massages (22), or orgasms (23), and consequently, OXT levels are higher in new lovers than in unattached singles (24). Nevertheless, it is currently unclear to what extent peripheral measurements reflect central OXT concentrations (25). An OXTR polymorphism has also been associated with pair-bonding traits in females (26). Intranasal administration of OXT increases ratings of facial attractiveness (27), positive communication behavior during couple conflicts (28), and may facilitate the fidelity of men in a monogamous relationship by making them keep a greater distance from other women (29). We therefore predicted that OXT would strengthen romantic bonds in men by increasing the perceived attractiveness and arousal evoked by face pictures of their female partner and concomitant neural activation in the VTA, NAcc, caudate, and lentiform nuclei.

To address this fascinating hypothesis, we conducted a discovery (DSC) and a replication (RPL) study, each involving a

Significance

Sexual monogamy is potentially costly for males, and few mammalian species along with humans exhibit it. The hypothalamic peptide oxytocin (OXT) has been implicated in mediating pair bonds in various species, but as yet, we know little about neurobiological factors that might act to promote fidelity, especially in men. Here we provide evidence for a mechanism by which OXT may contribute to romantic bonds in men by enhancing their partner's attractiveness and reward value compared with other women.

Author contributions: D.S., A.W., K.M.K., and R.H. designed research; D.S. and A.W. performed research; D.S., A.W., K.M.K., B.B., and R.H. analyzed data; B.S.-W. contributed new reagents/analytic tools; and D.S., A.W., K.M.K., B.S.-W., B.B., O.G., W.M., and R.H. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. A.S. is a guest editor invited by the Editorial Board.

¹D.S., A.W., and K.M.K. contributed equally to this work.

²To whom correspondence should be addressed. E-mail: renehurlemann@me.com.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10. 1073/pnas.1314190110/-/DCSupplemental.

counter balanced, double-blind, within-subject design experiment. In both studies, we used fMRI to repeatedly scan the brains of 20 heterosexual pair-bonded men (mean age ± SD and relationship length, DSC study: 25.1 ± 3.3 y and 28.8 ± 15.4 mo; RPL study: 26.6 ± 3.8 y and 36.4 ± 25.3 mo) after they received either intranasal OXT (24 IU) or placebo (PLC) treatment. During fMRI scanning of the DSC study, they were exposed to gray-scaled face photographs of their loved one, of an unfamiliar woman, or of a nonface control stimulus, a house. The photographs of other women were carefully matched for attractiveness, arousal, and picture quality with those of the partner in each case by using ratings obtained from a sample of 10 independent judges. Each picture of the three categories was presented 24 times (six blocks with four picture presentations). In the RPL study, we omitted the nonsocial control condition of the fMRI paradigm and instead used photographs of highly familiar, but nonrelated, women to control for possible nonspecific effects of OXT on familiarity. To avoid ceiling effects and to limit cognitive processing, we asked our subjects to passively view the pictures. After each fMRI scan, subjects completed the Passionate Love Scale (PLS) questionnaire (30) and rated the perceived attractiveness and arousal evoked by the faces using a visual analog scale ranging from 0 (minimum) to 100 (maximum). Further, love styles (e.g., Eros, a committed romantic relationship) (31) were measured before the administration of the nasal spray.

Results

All subjects reported being passionately in love (Table S1). A repeated-measures ANOVA with bond (DSC: partner vs. unfamiliar; RPL: partner, unfamiliar and familiar) and treatment (OXT vs. PLC) as within-subject variables and attractiveness ratings as a dependent variable yielded a main effect of bond (DSC: $F_{(1,19)} = 120.96$, P < 0.01, $\eta^2 = 0.88$; RPL: $F_{(2,18)} = 61.61$, P < 0.01, $\eta^2 = 0.87$) and an interaction effect of bond and treatment (DSC: $F_{(1,19)} = 7.00$, P = 0.02, $\eta^2 = 0.27$; RPL: $F_{(2,18)} = 2.82$, P = 0.086, $\eta^2 = 0.24$). For arousal ratings, we detected only a main effect of bond (DSC: $F_{(1,19)} = 74.96$, P < 0.01, $\eta^2 = 0.80$; RPL: $F_{(2,18)} = 84.47$, P < 0.01, $\eta^2 = 0.90$) and a main effect of

treatment in the DSC study ($F_{(1,19)} = 7.47, P = 0.01, \eta^2 = 0.28$), but no interaction effects (all P > 0.14). The main effect of bond became evident as a positive partner bias, i.e., the participants gave their partners significantly higher attractiveness and arousal ratings than other either unfamiliar or familiar women (attractiveness: DSC_{OXT}: $t_{(19)} = 12.11$, P < 0.01, d = 4.09; DSC_{PLC}: $t_{(19)} = 9.52, P < 0.01, d = 3.68; \text{RPL}_{\text{OXT}}: t_{(19)} = 10.92, P < 0.01,$ d = 3.90; RPL_{PLC}: $t_{(19)} = 9.68$, P < 0.01, d = 3.53 and arousal: DSC_{OXT}: $t_{(19)} = 11.03$, P < 0.01, d = 3.20; DSC_{PLC}: $t_{(19)} = 6.57$, $P < 0.01, d = 2.06; \text{RPL}_{\text{OXT}}: t_{(19)} = 12.21, P < 0.01, d = 3.51;$ RPL_{PLC}: $t_{(19)} = 8.87, P < 0.01, d = 2.60$). Interestingly, post hoc t tests in the RPL study also revealed a positive bias for attractiveness and arousal ratings of the highly familiar woman (attractiveness: RPL_{OXT}: $t_{(19)} = 8.12$, P < 0.01, d = 1.87; RPL_{PLC}: $t_{(19)} = 9.30, P < 0.01, d = 2.26$, arousal: RPL_{OXT}: $t_{(19)} = 8.12, P < 0.01$ 0.01, d = 2.14; RPL_{PLC}: $t_{(19)} = 9.30, P < 0.01, d = 2.13$), but these effects were significantly smaller than the partner bias (attractiveness: RPL_{OXT}: $t_{(19)} = 5.63$, P < 0.01, d = 1.39; RPL_{PLC}: $t_{(19)} =$ 5.79, P < 0.01, d = 1.17; arousal: RPL_{OXT}: $t_{(19)} = 4.44$, P < 0.01, d = 0.97; RPL_{PLC}: $t_{(19)} = 3.19$, P < 0.01, d = 0.84). The arousalbased treatment effect in the DSC study is related to higher arousal ratings under OXT, irrespective of the bond. However, this effect could not be replicated in the RPL study. Importantly, as indicated by the interaction effect of treatment and bond, OXT enhanced the positive partner bias in attractiveness ratings in both the DSC ($t_{(19)} = 2.65$, P = 0.02, d = 0.27; Fig. 1A) and RPL studies $(t_{(19)} = 2.32, P = 0.03, d = 0.17;$ Fig. 1C). OXT neither altered PLS scores nor the positive bias for familiar as opposed to unfamiliar women (all $\hat{P} > 0.05$). Moreover, a correlation analysis between the positive partner bias and love styles revealed that only a positive association with Eros was significant in the DSC and RPL studies, which survived Bonferroni correction for multiple testing (DSC: $r_{PLC} = 0.61$, P < 0.01; $r_{OXT} =$ 0.71, P < 0.01; Fig. 1B; RPL: $r_{PLC} = 0.79$, P < 0.01; $r_{OXT} = 0.73$, P < 0.01; Fig. 1D), providing further evidence that this bias indeed constitutes a fundamental element of a romantic relationship.

On the neural level, a whole-brain analysis in both the DSC and RPL study revealed an increased blood oxygen level-dependent (BOLD) signal in the middle occipital gyrus (DSC: MNI





Fig. 1. OXT effects on partner judgments. Both in the DSC (A) and RPL (C) study, the intranasal administration of OXT significantly increased the positive partner bias for attractiveness. This bias was positively correlated with Eros (a romantic love style) in the DSC (B) and RPL (D) study. Error bars indicate SEM. AR, arousal; AT, attractiveness; DSC, discovery; OXT, oxytocin; PLC, placebo; RPL, replication; *P < 0.05.

coordinates 30, -85, 4, $t_{(19)} = 6.85$, family wise error corrected: $P_{FWE} < 0.01$; RPL: MNI coordinates -39, -61, -8, $t_{(19)} = 5.74$, $P_{FWE} < 0.01$) and anterior cingulate cortex (DSC: MNI coordinates -6, 41, -5, $t_{(19)} = 4.46$, $P_{FWE} < 0.01$; RPL: MNI coordinates -3, 50, 7, $t_{(19)} = 5.74$, $P_{FWE} < 0.01$) when the partner's face was presented compared with an unfamiliar woman (Tables S2 and S3). We did not observe any comparable effects for the contrast [Partner > Familiar] and there were also no nonspecific main treatment effects across conditions (all P > 0.05).

To examine the specific OXT effect on the processing of the partner's face, we computed the contrast $[Partner_{OXT} >$ $Unfamiliar_{OXT}$ > [Partner_{PLC} > $Unfamiliar_{PLC}$] in both studies and additionally [Partner_{OXT} > Familiar_{OXT}] > [Partner_{PLC} > Familiar_{PLC}] in the RPL study. When we restricted our analysis to the VTA, NAcc, caudate, and lentiform nucleus as hypothesisdriven predefined regions of interest (ROIs), we found enhanced responsiveness for the contrast [Partner_{OXT} > Unfamiliar_{OXT}] > $[Partner_{PLC} > Unfamiliar_{PLC}]$ bilaterally in the NAcc (Fig. 2A) and VTA (Fig. 3A) ($P_{\rm FWE} < 0.05$; DSC study). This finding was replicated bilaterally in the NAcc (Fig. 2B) and unilaterally in the right VTA (Fig. 3B; RPL study). Inspection of the extracted individual percent signal changes confirmed that this effect was due to a differential response as a result of OXT treatment. With houses as control stimuli in the DSC study ([Partner_{OXT} > House_{OXT}] > $[Partner_{PLC} > House_{PLC}])$, we found similar OXT effects as with unfamiliar women for the NAcc (DSC left peak MNI coordinates $-6, 5, -5; t_{(114)} = 2.69, P_{FWE} = 0.03;$ right peak MNI coordinates 6, 2, -5; $t_{(114)} = 2.05$, $P_{\text{FWE}} = 0.098$). For the VTA, no significant effect was detected, indicating that not only an increased response to the partner but also a decreased response to alternative women is necessary for OXT effects to become evident. In fact, a direct comparison of the neural response to the unfamiliar woman under OXT and PLC ([Unfamiliar_{PLC} > Unfamiliar_{OXT}]) showed a trend for a reduced activation in the left VTA after OXT administration (DSC peak MNI coordinates -6, -10, -5; $t_{(19)} = 2.58$, $P_{\text{FWE}} =$ 0.051; Fig. S1).

To further elucidate the specificity of these findings, we applied the ROI analyses to the contrast [Partner_{OXT} > Familiar_{OXT}] > [Partner_{PLC} > Familiar_{PLC}] in the RPL study and again detected an increased activation in the NAcc (left peak MNI coordinates -9, 5, 1; $t_{(152)} = 2.73$, $P_{FWE} = 0.02$; right peak MNI coordinates 12, 5, 1; $t_{(152)} = 2.09$, $P_{FWE} = 0.099$; Fig. 44). Interestingly, OXT also enhanced the activation in response to the partner compared with the familiar control in the right putamen (peak MNI coordinates 27, -19, 7; $t_{(152)} = 3.65$, $P_{FWE} < 0.01$; Fig. 4B) and right globus pallidus (peak MNI coordinates 24, -7, -5; $t_{(152)} =$ 2.46, $P_{FWE} = 0.04$; Fig. 4C). For the contrast [Familiar_{OXT} > Unfamiliar_{OXT}] > [Familiar_{PLC} > Unfamiliar_{PLC}], there was no effect of OXT in either the NAcc or VTA, even at very liberal significance thresholds (uncorrected P < 0.05). However, we observed a greater activation in the caudate body (right: MNI x, y, z: 21, -13, 25, $t_{(152)} = 2.58$, $P_{FWE} = 0.03$), possibly suggesting that OXT may also influence nonpartner bonding via distinct striatal nuclei. As expected, neither the whole-brain nor the ROI-based analyses showed any OXT effect on responses to nonsocial control stimuli ([House_{OXT} > House_{PLC}] or [House_{OXT} < House_{PLC}]).

Discussion

In this study, we examined the effect of an intranasal OXT administration on the behavioral ratings of and the neural response to face stimuli of long-term romantic partners. Our results confirmed the hypothesis that OXT also contribute to romantic pair bonding in humans by showing that OXT treatment enhanced a positive bias in men to perceive their female partner's face as more attractive than that of other unfamiliar women, an effect paralleled by increased activation of reward neurocircuitry.

The notion that OXT may play a specific role in maintaining long-term pair bonding in humans is also substantiated by our previous findings that it makes men in a relationship keep a greater social distance from attractive female strangers (29) and reinforces the suggestion that OXT effects are both context and person dependent (32, 33). An OXT-mediated positive bias in the perception of the female partner can be adaptive and may serve as a long-term resilience factor because such idealization can predict sustained relationship satisfaction over the first 3 y of marriage (34). In the left NAcc, OXT augmented the neural response to the partner both in comparison with unfamiliar and familiar women. However, we also observed an increased neural response to nonpartner related bonding stimuli (familiar women) in the caudate body, intriguingly suggesting that OXT may influence different kinds of attachments via distinct striatal areas. Future studies are warranted to disentangle this complex interaction between OXT and different types of bonding. Importantly, three different phases in romantic relationships have been defined that



Fig. 2. OXT effects on NAcc responses. Both in the DSC (A) and RPL (B) study, the intranasal administration of OXT increased NAcc response to the female partner's face compared with a matched, but unfamiliar woman (OXT_(Partner > Unfamiliar) > PLC_(Partner > Unfamiliar); DSC: left peak MNI coordinates -6, 5, -5; $t_{(114)} = 3.22$, $P_{\text{FWE}} < 0.01$; right peak MNI coordinates 6, 2, -2; $t_{(114)} =$ 3.70, P_{FWE} < 0.01; RPL: left peak MNI coordinates -9, 5, 1; t₍₁₅₂₎ = 2.40, P_{FWE} = 0.052; right peak MNI coordinates 12, 2, 1; $t_{(152)} = 2.48$, $P_{FWE} = 0.04$; display threshold P < 0.05 uncorrected). Percent signal change in the bilateral NAcc showed the greatest response to the mate after OXT administration. Error bars indicate SEM. DSC, discovery; L, left hemisphere; NAcc, nucleus accumbens; OXT, oxytocin; PLC. placebo: R. right hemisphere: RPC, replication.



share common biological substrates but also involve different brain systems (11, 35, 36). The relative short first phase of "being in love" usually lasts around half a year and evolves into a longer phase of "passional love." Eventually, after several years, the third phase of a relationship can be characterized as "companionate love" similar to friendships. The mean relationship durations in the present study were 29 and 36 mo (DSC and RPL, respectively), indicating that the participants were in the second phase of their relationship in which the initial euphoria and excitation decline but the passion remains high. Thus, OXT may still play a substantial role in the process of establishing a pair bond in the first stage of romantic love. Attractiveness and arousal ratings may not be sufficiently sensitive to detect nonsexual bonding effects, but OXT also enhanced caudate responses to familiar control stimuli. Consequently, OXT may not exclusively affect long-term romantic partner- and parent-infant bonds (37), but also influence other kinds of social bonds.

Enhanced activation of the NAcc and VTA is likely to reflect an increased reward value of the female partner and suggests that OXT may affect these dopaminergic areas in the context of bonds in a similar manner to monogamous voles. Facial attractiveness is known to elicit striatal activation (38), but by choosing face stimuli matched for overall attractiveness, we avoided any nonspecific contribution from this. Likewise, our results cannot be accounted for by nonspecific conscious effects of OXT because it did not affect state anxiety or mood ratings, and the subjects could not identify whether they had received OXT or PLC (Tables S4 and S5). Furthermore, reduced VTA activation in response to the faces of other women after OXT treatment indicates it could further contribute to long-term pair bonding by making interactions with other women less rewarding, although we found no corresponding behavioral support for this. If the OXT effect on the VTA activation in response to the partner reflects a devaluation of other unfamiliar women the man has no social bond with, then this could explain why we did not find a similar response to other familiar women they do have a social bond with in the RPL study. Recently, it has been suggested that the VTA is the human brain site where oxytocin acts to attach salience to socially relevant cues (39), and it is also likely that the comparison of the partner with an unfamiliar woman involves a larger salience difference than the contrast with a familiar one.

The absence of a stronger neural response to the female partner during PLC treatment indicates that the presentation of

Fig. 3. OXT effects on VTA responses. Both in the DSC (A) and RPL (B) study, the intranasal administration of OXT increased neural responses in the VTA to the face of the female partner compared with a matched, but unfamiliar woman (OXT_(Partner > Unfamiliar) > PLC(Partner > Unfamiliar); DSC: left peak MNI coordinates -6, -10, -5; t₍₁₁₄₎ = 2.56, P_{FWE} = 0.03; right peak MNI coordinates 0, -10, -5; $t_{(114)} = 2.33$, $P_{FWE} = 0.051$; RPL: left peak MNI coordinates 0, -16, -8; $t_{(152)} =$ 1.93, $P_{FWE} = 0.12$; right peak MNI coordinates 6, -13, -5; $t_{(152)} = 2.89$, $P_{FWE} = 0.02$; display threshold P <0.05 uncorrected). Analyses of percent signal change in the bilateral VTA showed the greatest response to the mate after OXT administration. Error bars indicate SEM. DSC, discovery; L, left hemisphere; OXT, oxytocin; PLC, placebo; R, right hemisphere; RPC, replication; VTA, ventral tegmental area.

a partner's neutral expression face is not more rewarding per se than that of a novel female of equivalent attractiveness (40). Thus, findings in previous fMRI studies of increased activation in brain reward areas in response to a partner's face may reflect the instruction to think about the person at the same time (11– 14). In fact, another non-pharmaco-fMRI study in which no



Fig. 4. OXT effects on responses to the partner compared with a familiar woman. OXT increased the response to the female partner's face compared with a highly familiar woman in the left NAcc (*A*; peak MNI coordinates –9, 5, 1; $t_{(152)} = 2.73$, $P_{FWE} = 0.02$), right putamen (*B*; peak MNI coordinates 27, -19, 7; $t_{(152)} = 3.65$, $P_{FWE} < 0.01$), and right globus pallidus (*C*; peak MNI coordinates 24, -7, -5; $t_{(152)} = 2.46$, $P_{FWE} = 0.04$, display threshold P < 0.05 uncorrected). Error bars indicate SEM. DSC, discovery; L, left hemisphere; NAcc, nucleus accumbens; OXT, oxytocin; PLC, placebo; R, right hemisphere.

additional instruction was given also failed to detect any striatal activation in response to partner faces (41). Furthermore, previous studies differed with regard to sample characteristics (e.g., a higher mean relationship duration) and task design (e.g., a longer duration of stimulus presentation) (11–14).

For men in a relationship, increased endogenous OXT signaling in the brain following experience of proximity, social support, intimate contact, or sex with their romantic partner might make these behaviors even more rewarding via an engagement of related neurocircuitry. In this way, a feed-forward loop would be initialized, resulting in a progressive increase in desire for the partner, similar to a drug addict's increased craving for drug consumption (42). This notion suggests the intriguing possibility that OXT may serve as a potential treatment for love-related withdrawal syndromes, including lovesickness and pathological grief from loss of a loved one. However, against the background of mounting evidence for anxiogenic and fearenhancing OXT effects (33, 43, 44), it is necessary that the mechanisms determining the outcome of an OXT treatment are completely identified.

Although we report here a very specific OXT effect on pairbond-related behavioral ratings and neural substrates, several limitations of our study should be acknowledged. In accordance with the majority of previous intranasal OXT studies, we did not observe an OXT effect on state anxiety and mood ratings, but it is conceivable that more sensitive measures might have detected subtle alterations. Most importantly, we cannot rule out that lowlevel, unconscious changes in affective state occurred because we did not collect psychophysiological data. Furthermore, it is likely that social desirability contributed to the positive partner bias, and future studies should try to elucidate to what extent OXT influences the perception of the partner per se. However, the fact that OXT had no effect on the positive bias toward more socially desirable familiar as opposed to unfamiliar women indicates that social desirability alone cannot explain our behavioral findings.

Taken together, our findings suggest that OXT may contribute to romantic bonds by making men perceive their female partner as increasingly attractive and rewarding compared with other women. By keeping an exclusive devotion to a single mate, it may open an evolutionary path to biparental care of offspring and shift a potentially destructive competition between males for mates toward more beneficial in-group cooperation (45).

Materials and Methods

Participants. Twenty healthy, nonsmoking male adults participated in the DSC study, and a completely independent sample of 20 healthy, nonsmoking male adults volunteered in the RPL study after giving written informed consent. The studies were approved by the institutional review board of the Medical Faculty of the University of Bonn and were carried out in compliance with the latest revision of the Declaration of Helsinki. Subjects received monetary compensation for study participation. Screening of the subjects was conducted before the test sessions (Table S6). Subjects were free of current and past physical or psychiatric illness, as assessed by medical history and a structured clinical interview for axis I and axis II disorders (DSM) IV (46). In addition, they were naive to prescription-strength psychoactive medication and had not taken any over-the-counter psychoactive medication in the last 4 wk.

All subjects were in a romantic heterosexual relationship for more than 6 mo, were unmarried, and had no children. The subjects reported to be passionately in love, and the time intervals since they last saw their partners and had intimate contact were comparable between the OXT and PLC sessions, as well as between the DSC and RPL study (Table 51). We also controlled if the subjects had an argument with their partners in the week before both test sessions and if anything important in their relationship changed between the two test sessions. The familiar control in the RPL study was a nonrelated woman whom the participants had known for at least 30 mo.

Experimental Design. In both the DSC and RPL studies, we applied a randomized, placebo-controlled, double-blind, within-group design. Subjects were randomly assigned to either intranasal administration of OXT (24 IU; Syntocinon-Spray; Novartis; three puffs per nostril, each with 4 IU OXT) or PLC (sodium chloride solution) 30 min before the start of the fMRI. The mean interval between the two fMRI sessions was 33 d (minimum, 21 d; maximum, 56 d) in the DSC study and 12.7 d (minimum, 6 d; maximum, 35 d) in the RPL study. Details on the tasks, fMRI procedure, and analyses can be found in *SI Materials and Methods*.

fMRI Paradigm. We adopted a modified version of an established face perception task (47) in the DSC study involving passive viewing of photographs of the partner, a matched, unknown woman, and houses. In the RPL study, we used photographs of the partner and of a highly familiar woman. As control stimuli, a photograph of an unknown woman matched to the partner and another photograph matched to the familiar woman were used in the RPL study. Subjects were asked to send several photographs of their partners and of a familiar, but nonrelated, woman, arranged as passport photographs, with a neutral, but friendly facial expression. The selected photographs were equated for facial expression, size, and luminance. An independent sample of 10 heterosexual men (age: 25.50 \pm 2.99 y; an ANOVA with the factor group yielded no significant age difference between the DSC, RPL, or pilot samples, P = 0.38) rated the arousal and attractiveness of the women and the quality of the photographs on a visual analog scale (0 = minimum, 100 = maximum) before the first fMRI session. Based on these three dimensions, the Euclidian distances between all photographs were computed, and the photograph with the lowest Euclidian distance was chosen as control stimulus for a partner photograph. The same procedure was applied to find matched photographs for the familiar women in the RPL study. We ensured that no subject knew the woman whose photograph was used as his control stimulus. To avoid confounding factors in the photographs, the backgrounds were masked in black, and the photographs were grav-scaled. Pictures of houses were used as nonfacial control stimuli. Houses serve as an appropriate control stimuli for faces because they share similar spatial and visual features (48). Moreover, houses are everyday objects with nonarousing properties. All houses were gray-scaled and equated for size and luminance.

Using Presentation 14 (Neurobehavioral Systems), stimuli were presented blockwise, via liquid crystal display (LCD) video goggles (Nordic NeuroLab). In total, there were six blocks for each stimulus category, and each block included four repetitions of the same stimulus (e.g., four times the photograph of the partner). Stimuli were presented for 2,625 ms on-screen, and the interstimulus interval varied between 250 and 1,500 ms to create jitter, resulting in a mean block length of 14.5 s. The sequence of blocks was randomized, and blocks were separated from each other by a low-level baseline period of 14.5-s duration, where a fixation cross was depicted in the center of the screen. To assure attentive stimulus was presented (percent correct responses DSC: OXT 97.64 \pm 7.49, PLC 95.90 \pm 10.16; RPL: OXT 98.62 \pm 3.19, PLC 98.62 \pm 4.5; all $P \geq 0.33$). In addition, the subjects were asked to look attentively at the pictures and to avoid cognitive processing; there were no further instructions.

Behavioral Task. After scanning, subjects were seated in front of a computer and were asked to rate photographs showing their own partner, the familiar woman, and partners of the other subjects on a visual analog scale for attractiveness (ranging from 0, most unattractive, to 100, most attractive) and arousal (ranging from 0, not arousing, to 100, most arousing). The sequence of the photographs was randomized.

Acquisition of fMRI Data. The MRI data of the DSC and RPL study were acquired with the same system and scanning parameters. A Siemens Trio MRI system (Siemens) operating at 3 T was used to obtain T2*-weighted echoplanar (EPI) images with BOLD contrast (repetition time = 3,000 ms, echo time = 35 ms, matrix size: 64×64 , pixel size: $3 \times 3 \times 3$ mm, slice thickness = 3.0 mm, distance factor = 10%, field of view = 192, flip angle = 90°, 36 axial slices). In addition, high-resolution anatomical images were acquired on the same scanner using a T1-weighted 3D MPRAGE sequence (imaging parameters: repetition time = 1,570 ms, echo time = 3.42 ms, matrix size: 256×256 , pixel size: $1 \times 1 \times 1$ mm, slice thickness = 1.0 mm, field of view = 256, flip angle = 15°, 160 sagital slices).

Analysis of fMRI Data. fMRI data were preprocessed and analyzed using SPM8 software (Wellcome Trust Centre for Neuroimaging, www.fil.ion.ucl.ac.uk/ spm) implemented in Matlab 7 (MathWorks). On the first level, six conditions (Partner_{OXT}, Control_{OXT}, House_{OXT}, Partner_{PLC}, Control_{PLC}, and House_{PLC}) in the DSC study and eight conditions (Partner_{OXT}, Control_Partner_{OXT}, Familiar_{OXT}, Control_Familiar_{OXT}, Partner_{PLC}, Control_Partner_{PLC}, Familiar_{PLC}, and

Control_Familiar_{PLC}) in the RPL study were modeled by a boxcar function convolved with a hemodynamic response function (49). The movement parameters were included as confounds in the design matrix. Main effects of treatment and bonding were analyzed by comparing the conditions relative to the low-level baseline. Parameter estimates for each contrast were subjected to one-sample *t* tests on the second level for the whole brain with a significance threshold of *P* < 0.05 corrected for multiple comparisons [familywise error (FWE)]. To specifically examine the modulatory effects of OXT, a flexible factorial model was designed with the factors treatment (OXT vs. PLC) and bond (DSC: partner vs. unfamiliar; RPL: partner, unfamiliar and familiar) to test the contrasts [Partner_{OXT} > Control_{OXT}] > [Partner_{PLC} > Control_{PLC}], [Partner_{OXT} > House_{OXT}] > [Partner_{PLC} > House_{PLC}], [Familiar_{OXT} >

- 1. Kleiman DG (1977) Monogamy in mammals. Q Rev Biol 52(1):39-69.
- Young KA, Gobrogge KL, Liu Y, Wang Z (2011) The neurobiology of pair bonding: Insights from a socially monogamous rodent. Front Neuroendocrinol 32(1):53–69.
- Smeltzer MD, Curtis JT, Aragona BJ, Wang Z (2006) Dopamine, oxytocin, and vasopressin receptor binding in the medial prefrontal cortex of monogamous and promiscuous voles. *Neurosci Lett* 394(2):146–151.
- Shapiro LE, Insel TR (1992) Oxytocin receptor distribution reflects social organization in monogamous and polygamous voles. Ann N Y Acad Sci 652:448–451.
- Ophir AG, Gessel A, Zheng DJ, Phelps SM (2012) Oxytocin receptor density is associated with male mating tactics and social monogamy. *Horm Behav* 61(3):445–453.
- Aragona BJ, et al. (2006) Nucleus accumbens dopamine differentially mediates the formation and maintenance of monogamous pair bonds. *Nat Neurosci* 9(1):133–139.
- Liu Y, Wang ZX (2003) Nucleus accumbens oxytocin and dopamine interact to regulate pair bond formation in female prairie voles. *Neuroscience* 121(3):537–544.
- Liu Y, Curtis JT, Wang Z (2001) Vasopressin in the lateral septum regulates pair bond formation in male prairie voles (Microtus ochrogaster). *Behav Neurosci* 115(4):910–919.
- Lim MM, Young LJ (2004) Vasopressin-dependent neural circuits underlying pair bond formation in the monogamous prairie vole. *Neuroscience* 125(1):35–45.
- Cho MM, DeVries AC, Williams JR, Carter CS (1999) The effects of oxytocin and vasopressin on partner preferences in male and female prairie voles (Microtus ochrogaster). *Behav Neurosci* 113(5):1071–1079.
- Acevedo BP, Aron A, Fisher HE, Brown LL (2012) Neural correlates of long-term intense romantic love. Soc Cogn Affect Neurosci 7(2):145–159.
- Aron A, et al. (2005) Reward, motivation, and emotion systems associated with earlystage intense romantic love. J Neurophysiol 94(1):327–337.
- Bartels A, Zeki S (2000) The neural basis of romantic love. Neuroreport 11(17): 3829–3834.
- Bartels A, Zeki S (2004) The neural correlates of maternal and romantic love. Neuroimage 21(3):1155–1166.
- Love TM, et al. (2012) Oxytocin gene polymorphisms influence human dopaminergic function in a sex-dependent manner. *Biol Psychiatry* 72(3):198–206.
- Romero-Fernandez W, Borroto-Escuela DO, Agnati LF, Fuxe K (2013) Evidence for the existence of dopamine d2-oxytocin receptor heteromers in the ventral and dorsal striatum with facilitatory receptor-receptor interactions. *Mol Psychiatry* 18(8):849–850.
- 17. Insel TR (2010) The challenge of translation in social neuroscience: A review of oxytocin, vasopressin, and affiliative behavior. *Neuron* 65(6):768–779.
- Hurlemann R, et al. (2010) Oxytocin enhances amygdala-dependent, socially reinforced learning and emotional empathy in humans. J Neurosci 30(14):4999–5007.
- Striepens N, Kendrick KM, Maier W, Hurlemann R (2011) Prosocial effects of oxytocin and clinical evidence for its therapeutic potential. Front Neuroendocrinol 32(4):426–450.
- Light KC, Grewen KM, Amico JA (2005) More frequent partner hugs and higher oxytocin levels are linked to lower blood pressure and heart rate in premenopausal women. *Biol Psychol* 69(1):5–21.
- Grewen KM, Girdler SS, Amico J, Light KC (2005) Effects of partner support on resting oxytocin, cortisol, norepinephrine, and blood pressure before and after warm partner contact. *Psychosom Med* 67(4):531–538.
- Morhenn V, Beavin LE, Zak PJ (2012) Massage increases oxytocin and reduces adrenocorticotropin hormone in humans. *Altern Ther Health Med* 18(6):11–18.
- 23. Borrow AP, Cameron NM (2012) The role of oxytocin in mating and pregnancy. Horm Behav 61(3):266–276.
- Schneiderman I, Zagoory-Sharon O, Leckman JF, Feldman R (2012) Oxytocin during the initial stages of romantic attachment: Relations to couples' interactive reciprocity. *Psychoneuroendocrinology* 37(8):1277–1285.

Control_Familiar_{OXT}] > [Familiar_{PLC} > Control_Familiar_{PLC}], and [Partner_{OXT} > Familiar_{OXT}] > [Partner_{PLC} > Familiar_{PLC}]. The Wake Forest University Pickatlas (Version 3.0) was used to generate 5-mm ROI masks for NAcc, VTA, caudate, and lentiform nucleus (*SI Materials and Methods*), and the threshold for significance was set at P < 0.05 and was FWE corrected for multiple comparisons based on the size of the ROI.

ACKNOWLEDGMENTS. R.H. was supported by German Research Foundation Grant HU1302/2-2 and a starting independent researcher grant (Neuromodulation of Emotion) jointly provided by the Ministry of Innovation, Science, Research and Technology of the German State of North Rhine-Westphalia and the University of Bonn. K.M.K. was supported by National Natural Science Foundation of China Grant 91132720.

- Churchland PS, Winkielman P (2012) Modulating social behavior with oxytocin: How does it work? What does it mean? *Horm Behav* 61(3):392–399.
- Walum H, et al. (2012) Variation in the oxytocin receptor gene is associated with pairbonding and social behavior. *Biol Psychiatry* 71(5):419–426.
- Theodoridou A, Rowe AC, Penton-Voak IS, Rogers PJ (2009) Oxytocin and social perception: Oxytocin increases perceived facial trustworthiness and attractiveness. *Horm Behav* 56(1):128–132.
- Ditzen B, et al. (2009) Intranasal oxytocin increases positive communication and reduces cortisol levels during couple conflict. *Biol Psychiatry* 65(9):728–731.
- Scheele D, et al. (2012) Oxytocin modulates social distance between males and females. J Neurosci 32(46):16074–16079.
- Hatfield E, Sprecher S (1986) Measuring passionate love in intimate relationships. J Adolesc 9(4):383–410.
- Lee JA (1973) Colours of Love: An Exploration of the Ways of Loving (New Press, Toronto).
- Bartz JA, Zaki J, Bolger N, Ochsner KN (2011) Social effects of oxytocin in humans: Context and person matter. *Trends Cogn Sci* 15(7):301–309.
- Striepens N, et al. (2012) Oxytocin facilitates protective responses to aversive social stimuli in males. Proc Natl Acad Sci USA 109(44):18144–18149.
- Murray SL, et al. (2011) Tempting fate or inviting happiness?: Unrealistic idealization prevents the decline of marital satisfaction. *Psychol Sci* 22(5):619–626.
- de Boer A, van Buel EM, Ter Horst GJ (2012) Love is more than just a kiss: A neurobiological perspective on love and affection. *Neuroscience* 201:114–124.
- Fisher HE, Aron A, Brown LL (2006) Romantic love: A mammalian brain system for mate choice. *Philos Trans R Soc Lond B Biol Sci* 361(1476):2173–2186.
- Feldman R (2012) Oxytocin and social affiliation in humans. Horm Behav 61(3): 380–391.
- Aharon I, et al. (2001) Beautiful faces have variable reward value: fMRI and behavioral evidence. *Neuron* 32(3):537–551.
- 39. Groppe SE, et al. (2013) Oxytocin influences processing of socially relevant cues in the ventral tegmental area of the human brain. *Biol Psychiatry* 74(3):172–179.
- Bunzeck N, Düzel E (2006) Absolute coding of stimulus novelty in the human substantia nigra/VTA. Neuron 51(3):369–379.
- Taylor MJ, et al. (2009) Neural correlates of personally familiar faces: Parents, partner and own faces. Hum Brain Mapp 30(7):2008–2020.
- Burkett JP, Young LJ (2012) The behavioral, anatomical and pharmacological parallels between social attachment, love and addiction. *Psychopharmacology (Berl)* 224(1): 1–26.
- Guzmán YF, et al. (2013) Fear-enhancing effects of septal oxytocin receptors. Nat Neurosci 16(9):1185–1187.
- Grillon C, et al. (2013) Oxytocin increases anxiety to unpredictable threat. Mol Psychiatry 18(9):958–960.
- Gavrilets S (2012) Human origins and the transition from promiscuity to pair-bonding. Proc Natl Acad Sci USA 109(25):9923–9928.
- American Psychiatric Association (2000) Diagnostic and Statistical Manual of Mental Disorders (American Psychiatric Association, Washington), 4th Ed.
- 47. Goossens L, et al. (2009) Selective processing of social stimuli in the superficial amygdala. *Hum Brain Mapp* 30(10):3332–3338.
- Yovel G, Kanwisher N (2004) Face perception: Domain specific, not process specific. Neuron 44(5):889–898.
- Friston KJ, et al. (1995) Statistical parametric maps in functional imaging: A general linear approach. Hum Brain Mapp 2(4):189–210.

Supporting Information

Scheele et al. 10.1073/pnas.1314190110

SI Materials and Methods

Study Subjects. All subjects completed a comprehensive neuropsychological test battery. In the discovery (DSC) study, neuropsychological testing before study enrollment included the German version of the Rey Auditory Verbal Learning Test (RAVLT) (1, 2) to assess verbal learning skills, the digit-span test (DST) derived from the revised Wechsler adult intelligence scale (3) to assess working memory performance, the Leistungspruefsystem Subtest 4 (LPS 4) (4) to assess nonverbal reasoning IQ, the Mehrfach-Wortschatz-Intelligenztest Teil B (MWT-B) (5) to assess verbal IQ based on lexical decisions, and the trailmaking test (TMT) (6) parts A and B to assess visual attention and task-switching performance. In the replication (RPL) study, cognitive performance was assessed using the Cambridge Neuropsychological Test Automated Battery (CANTAB), a computerized neurocognitive assessment presented through a touchscreen computer (7). For details of the outcome measure, see the CANTABeclipse Test Administration Guide (8). Subjects' speed of response to a visual target, the ability to retain spatial information, and visual memory were measured with the simple and reaction time task (RTI), the spatial working memory task (SWM; eight boxes version), and the paired associates learning task (PAL), respectively. All subjects were within a normal range of cognitive performance (Table S6).

The participants were asked to maintain their regular bed and waking times and to abstain from caffeine and alcohol intake on the day of the experiment. To control for potentially confounding effects of oxytocin (OXT) on state anxiety and mood, all subjects completed the State-Trait Anxiety Inventory (STAI) (9) and the Positive and Negative Affective Scale (PANAS) (10) immediately before the OXT/placebo administration and after the experimental task. Furthermore, all subjects completed the d2 Test of Attention (Aufmerksamkeits- und Belastungstest d2) (11) after the experimental task. For both the DSC and RPL study data, three repeated-measure ANOVAs with measurement (before and after the experiment) and treatment (OXT and placebo) as within-subject factors and state anxiety, positive affect, or negative affect as dependent variables revealed no significant main or interaction effects (all P > 0.12). There was also no significant difference between the d2 attention performance of the OXT and placebo (PLC) sessions in the DSC and RPL study (all P > 0.06; cf. Tables S4 and S5). Thus, OXT did not influence subjective anxiety, mood ratings, or attention. After completing the task, subjects were debriefed and asked to guess whether they had received OXT or PLC. The estimation of the received treatment was comparable between the OXT and PLC session in the DSC (correct estimates: OXT, n = 11; PLC, n = 9; $\chi^{2}_{(1)} =$ 0.9; P = 0.34) and RPL study (correct estimates: OXT, n = 9; PLC, n = 8; $\chi^2_{(1)} = 0.12$; P = 0.73), showing that the subjects were unaware of whether they had received OXT or PLC.

Functional MRI Paradigm. An independent sample of 10 heterosexual healthy men (mean age \pm SD: 25.50 \pm 2.99 y; an ANOVA with the factor group yielded no significant age difference between the DSC, RPL, or pilot samples, P = 0.38) rated attractiveness of and the arousal induced by partner, familiar, and unfamiliar women, as well as the quality of the photographs on a visual analog scale (0 = minimum, 100 = maximum) before the first functional MRI (fMRI) session. The ratings of each dimension were adequately intercorrelated (DSC: estimated correlation of arousal ratings, $\rho = 0.63$; attractiveness, $\rho = 0.88$; picture quality, $\rho = 0.90$; RPL: arousal, $\rho = 0.56$; attractiveness, $\rho = 0.85$; picture quality, $\rho = 0.93$). In total, the task lasted around 10 min in the DSC study and 12 min (due to the additional trials with the matched controls for highly familiar women) in the RPL study.

Behavioral Task. Before the start of the experiment, the subjects were asked if they knew any of the control persons, and ratings of familiar persons (except for the partner and the familiar woman in the RPL study) were discarded from further analysis (DSC: 3 ratings, RPL: 81 ratings). After the ratings, all subjects completed the Passionate Love Scale (PLS) (12), which had a good internal consistency in both sessions (DSC: OXT Cronbach's $\alpha = 0.85$, PLC $\alpha = 0.91$; RPL: OXT $\alpha = 0.93$, PLC $\alpha = 0.92$). A 1 (minimum) to 9 (maximum) scale was used for the PLS. Example items are "I want _____ to know me-my thoughts, my fears, and my hopes" or "Sometimes I can't control my thoughts; they are obsessively on _____." Furthermore, all subjects completed the Marburg Attitude Scales towards Love Styles (MEIL), which is a German version of love styles developed by Lee (13). It contains three primary styles of loving: the first one is Eros, a romantic love style that is similar to passionate love and is characterized by a powerful attraction to the beloved individual. Interestingly, a dopamine D2 receptor (D2R) polymorphism has been found to be associated with the Eros love scale (14). The second is Ludus, which describes lovers who view love as a game and often have several partners simultaneously. The third is Storge, a slow developing, friendship-based love. These primary love styles can be combined to form secondary styles of love: Pragma (Storge and Ludus combined; pragmatic view on the relationship), Mania (Eros and Ludus combined; obsessive and possessive lover), and Agape (Storge and Eros combined; altruistic love style). In the German version, each love style is assessed with 10 items, and each dimension has an adequate reliability (DSC: Eros $\alpha = 0.86$, Ludus $\alpha = 0.88$, Storge $\alpha = 0.73$, Pragma $\alpha = 0.74$, Mania $\alpha = 0.88$, Agape $\alpha = 0.74$; RPL: Eros $\alpha = 0.94$, Ludus $\alpha = 0.48$, Storge $\alpha = 0.74$, Pragma $\alpha = 0.80$, Mania $\alpha = 0.77$, Agape $\alpha = 0.88$). For the samples of the DSC and RPL study, we observed the highest scores for Eros and the lowest scores for Ludus (cf. Table S1).

Analysis of fMRI Data. The first five volumes of each functional time series were discarded to allow for T1 equilibration. Images were corrected for head movement between scans by an affine registration. For realignment, a two-pass procedure was used, by which images were initially realigned to the first image of the time series and subsequently rerealigned to the mean of all images. For spatial normalization, the mean EPI image of each subject was normalized to the current Montreal Neurological Institute (MNI) template (15, 16) using the unified segmentation function in SPM-8. This algorithm combines image registration, tissue classification, and bias correction within the same generative model. All images were thereby transformed into standard stereotaxic space and resampled at $3 \times 3 \times 3$ -mm voxel size. The normalized images were spatially smoothed using an 8-mm full width at half maximum Gaussian kernel. Raw time series were detrended by the application of a high-pass filter (cutoff period, 128 s). A twolevel random effects approach based on the general linear model as implemented in SPM-8 was used for statistical analyses.

Based on previous studies investigating the neural correlates of romantic love (17–19), we used 5-mm spheres as regions of interest (ROIs) centered at the coordinates of the reported maximum value for the nucleus accumbens (left: -10, 4, -4; right: 10, 4, -4), ventral tegmental area (left: -2, -12, -8; right: 2,

-12, -8), caudate body (left: -18, -14, 22; right: 18, -14, 22), caudate tail (left: -34, -32, -4; right: 36, -34, 0), putamen (left: -22, 2, 4; right: 24, -18, 10), and globus pallidus posterior (left: -32, 6, -8; right: 24, -8, -8). The Wake Forest University (WFU) Pickatlas (version 3.0) was used to generate ROI masks, and the threshold for significance was set at P < 0.05 and familywise error (FWE) corrected for multiple comparisons based on the size of the ROI.

Salivary OXT Collection and Analysis. Saliva samples were collected using prechilled Salivettes (Sarstedt). One sample was collected before administration of the nasal spray both in the OXT and PLC session and another sample was collected after the fMRI task. Salivettes were immediately centrifuged at 4,180 \times g for 2 min, and aliquoted samples were stored at -80 °C until assayed. Salivary OXT concentrations were determined by using a 96-well commercial OXT-ELISA kit (ENZO). Measurements were performed in duplicate, and samples were treated according to kit instructions. According to the manufacturer, the sensitivity limit of the assay is 11.7 pg/mL, and 15.1% of the samples fell below the lower level of sensitivity. The assay's reported intraassay and interassay coefficients of variability are 9.1–12.4% and 5.2–14.5%, respectively.

Statistical Analysis. Demographical, neuropsychological, and behavioral data were analyzed using SPSS 20 (SPSS Inc.). Quantitative behavioral data were compared by repeated-measures ANOVA, and Pearson's product-moment correlation was used for correlation analysis. Eta-squared and Cohen's *d* were calculated as measures of effect size. The assumption of normality for all target variables was assessed separately for the OXT and PLC sessions using Kolmogorov-Smirnov tests. All target data were derived from normally distributed populations (all P > 0.06). The assumption of sphericity was assessed with Mauchly's test, and for significant violations, Greenhouse-Geisser's correction was applied. For qualitative variables, Pearson's χ^2 tests were used. All reported *P* values are two-tailed, if not otherwise noted, and P < 0.05 was considered significant.

SI Results

Behavioral Results. In both the DSC and RPL study, the OXT effect on the positive partner bias was dependent on a higher perceived attractiveness of the partner (DSC: $t_{(19)} = 1.75$, P = 0.048 one-tailed, d = 0.27; RPL: $t_{(19)} = 1.64$, P = 0.059 one-tailed, d = 0.16) rather than on a derogation of other women (DSC: $t_{(19)} = -1.12$, P = 0.28, d = -0.14; RPL: $t_{(19)} = -0.78$, P = 0.44, d = -0.12).

A slightly different definition of the positive partner bias yielded similar results. The positive partner bias defined as the difference between the attractiveness of the partner rated by the partner or other participants was significantly larger in the OXT (DSC: 41.37 ± 15.28 ; RPL: 44.89 ± 14.06) than in the PLC session

- 1. Rey LB (1941) L'examen psychologique dans les cas d'encephalopathie traumatique. Archives of Psychology 28:215–285.
- Helmstaedter C, Lendt M, Lux S (2001) VLMT—Verbaler Lern—und Merkfähigkeitstest (Beltz Test GmbH, Göttingen, Germany).
- 3. Wechsler D (1997) Wechsler Adult Intelligence Scale. Administration and Scoring Manual (Psychological Corporation, San Antonio, TX), 3rd Ed.
- Horn W (1983) Leistungsprüfsystem L-P-S (Hogrefe, Göttingen, Germany).
 Lehrl S, Triebig G, Fischer B (1995) Multiple choice vocabulary test MWT as a valid and
- Lenn's, Triebig G, Fischer B (1995) Multiple choice vocabulary test MWI as a valid and short test to estimate premorbid intelligence. Acta Neurol Scand 91(5):335–345.
- Raitan RM (1958) Validity of the trail making test as an indication of organic brain damage. Percept Mot Skills 8:271–276.
- Sahakian BJ, Owen AM (1992) Computerized assessment in neuropsychiatry using CANTAB: discussion paper. J R Soc Med 85(7):399–402.
- 8. CANTABeclipse (2011) Test Administration Guide (Cambridge Cognition Limited, Cambridge, UK).
- 9. Spielberger CD, Gorsuch RL, Lushene RE (1970) Manual for the State-Trait Anxiety Inventory (Consulting Psychologists Press, Palo Alto, CA).

(DSC: 36.50 ± 13.89 ; RPL: 41.25 ± 13.50) (DSC: $t_{(19)} = 2.46$, P = 0.02, d = 0.33; RPL: $t_{(19)} = 2.53$, P = 0.02, d = 0.26).

Salivary OXT Concentrations. In the RPL study, we also measured salivary OXT concentrations before the nasal spray administration (pre) and again after the fMRI task (post) to examine potential changes in the endogenous OXT levels in the PLC session due to the repeated presentation of partner and familiar woman photographs. In the OXT session, the concentration rose dramatically (pre: 34 ± 24 pg/mL, post: 589 ± 1022 pg/mL, Z = 3.92, P < 0.01), which is consistent with previous studies (20) but may also partly be attributed to OXT leaking from the nasal cavity into the mouth. More importantly, after the fMRI task, the endogenous OXT levels were also increased in the PLC session (pre: 35 ± 24 pg/mL, post: 57 ± 51 pg/mL, Z = 2.58, P = 0.01), which may indicate that face pictures of the partner and familiar individuals facilitated endogenous OXT release in line with a report in sheep (21), although we cannot rule out an effect of stress caused by the noisy scanner environment. Peripheral saliva samples have been used in previous studies (20, 22, 23), but the validity of saliva OXT measurement for quantification purposes has been questioned (24, 25), and the association between peripheral and central OXT level in the brain is highly controversial (26–28). Thus, these data should be interpreted as reflecting relative change rather than indicating absolute quantities.

fMRI Results. We found no significant association between the neural response in the nucleus accumbens (NAcc) and ventral tegmental area (VTA) and the OXT-induced changes in attractiveness ratings. This absence of correlation could be related to the absence of behavioral ratings performed during the fMRI scan, but it is also conceivable that the behavioral and neural OXT effects in this study are independent of each other.

Importantly, we also found no evidence for unspecific nonsocial OXT effects because there was no significant effect for the contrast [House_{OXT} > House_{PLC}] in any ROI, even at a very low significance threshold (P < 0.001 uncorrected, extent cluster threshold of $k \ge 0$ voxels). The results for the contrast [Partner > Unfamiliar] are shown in Tables S2 and S3 separately for the OXT and PLC sessions.

Furthermore, we were interested in deactivation in response to images of the partner. For the DSC and RPL study, on the whole-brain level, we found no significant deactivation for the contrasts comparing the OXT and PLC sessions ([Unfamiliar_{PLC} > Partner_{PLC}] > [Unfamiliar_{OXT} > Partner_{OXT}], [House_{PLC} > Partner_{PLC}] > [House_{OXT} > Partner_{OXT}], [House_{PLC} > Partner_{PLC}]). Only in the PLC session of the RPL study, but not in the DSC study, did the contrast [Unfamiliar_{PLC} > Partner_{PLC}] yield a significant cluster in the post-central gyrus (MNI *x*, *y*, *z*: -45, -22, 58; $t_{(19)} = 4.70$; $P_{FWE} = 0.004$).

- Watson D, Clark LA, Tellegen A (1988) Development and validation of brief measures of positive and negative affect: The PANAS scales. J Pers Soc Psychol 54(6):1063–1070.
- 11. Brickenkamp R, Zillmer E (1998) The d2 Test of Attention (Hogrefe & Huber Publishers, Seattle), 1st Ed.
- Hatfield E, Sprecher S (1986) Measuring passionate love in intimate relationships. J Adolesc 9(4):383–410.
- Lee JA (1988) Love styles. The Psychology of Love, eds Barnes MH, Sternberg RJ (Yale Univ Press, New Haven, CT), pp 38–67.
- Emanuele E, Brondino N, Pesenti S, Re S, Geroldi D (2007) Genetic loading on human loving styles. Neuroendocrinol Lett 28(6):815–821.
- Evans AC, et al. (1992) Anatomical mapping of functional activation in stereotactic coordinate space. *Neuroimage* 1(1):43–53.
- Holmes CJ, et al. (1998) Enhancement of MR images using registration for signal averaging. J Comput Assist Tomogr 22(2):324–333.
- 17. Bartels A, Zeki S (2000) The neural basis of romantic love. *Neuroreport* 11(17): 3829–3834.

- Bartels A, Zeki S (2004) The neural correlates of maternal and romantic love. Neuroimage 21(3):1155–1166.
- Acevedo BP, Aron A, Fisher HE, Brown LL (2012) Neural correlates of long-term intense romantic love. Soc Cogn Affect Neurosci 7(2):145–159.
- Weisman O, Zagoory-Sharon O, Feldman R (2012) Intranasal oxytocin administration is reflected in human saliva. *Psychoneuroendocrinology* 37(9):1582–1586.
- Fabre-Nys C, Ohkura S, Kendrick KM (1997) Male faces and odours evoke differential patterns of neurochemical release in the mediobasal hypothalamus of the ewe during oestrus: An insight into sexual motivation? *Eur J Neurosci* 9(8):1666–1677.
- Huffmeijer R, et al. (2012) Salivary levels of oxytocin remain elevated for more than two hours after intranasal oxytocin administration. Neuroendocrinol Lett 33(1):21–25.
- van Ijzendoorn MH, Bhandari R, van der Veen R, Grewen KM, Bakermans-Kranenburg MJ (2012) Elevated Salivary Levels of Oxytocin Persist More than 7 h after Intranasal Administration. Front Neurosci 6:174.
- Horvat-Gordon M, Granger DA, Schwartz EB, Nelson VJ, Kivlighan KT (2005) Oxytocin is not a valid biomarker when measured in saliva by immunoassay. *Physiol Behav* 84(3):445–448.
- Carter CS, et al. (2007) Oxytocin: Behavioral associations and potential as a salivary biomarker. Ann N Y Acad Sci 1098:312–322.
- Kagerbauer SM, et al. (2013) Plasma oxytocin and vasopressin do not predict neuropeptide concentrations in human cerebrospinal fluid. J Neuroendocrinol 25(7): 668–673.
- 27. Wang YL, et al. (2013) The interaction between the oxytocin and pain modulation in headache patients. *Neuropeptides* 47(2):93–97.
- McCullough ME, Churchland PS, Mendez AJ (2013) Problems with measuring peripheral oxytocin: Can the data on oxytocin and human behavior be trusted? *Neurosci Biobehav Rev* 37(8):1485–1492.



Fig. S1. OXT effects on NAcc responses with houses as control stimuli. The intranasal administration of OXT increased NAcc response to the female partner's face compared with houses ($OXT_{(Partner > House)} > PLC_{(Partner > House)}$; DSC left peak MNI coordinates -6, 5, -5; $t_{(114)} = 2.69$, $P_{FWE} = 0.03$; right peak MNI coordinates 6, 2, -5; $t_{(114)} = 2.05$, $P_{FWE} = 0.098$; display threshold P < 0.05 uncorrected). Percent signal change in the bilateral NAcc showed the greatest response to the mate after OXT administration. Error bars indicate SEM. DSC, discovery; L, left hemisphere; NAcc, nucleus accumbens; OXT, oxytocin; PLC, placebo; R, right hemisphere.

Variable	DSC study [mean (±SD)]	RPL study [mean (±SD)]
Relationship duration (months)	28.75 (15.43)	36.35 (25.33)
Age of partner (years)	24.00 (3.87)	24.90 (3.64)
PLS OXT* ^{,†}	6.33 (0.96)	6.46 (1.36)
PLS PLC* ^{,†}	6.46 (1.21)	6.49 (1.27)
Time (d) since the last time seen OXT*	1.75 (4.52)	1.84 (4.06)
Time (d) since the last time seen PLC*	0.60 (1.27)	1.05 (2.53)
Time (d) since the last intimate contact OXT*	2.55 (4.41)	4.26 (4.31)
Time (d) since the last intimate contact PLC*	3.95 (6.83)	3.32 (3.02)
Love style Eros [‡]	7.19 (0.99)	6.98 (1.55)
Love style Ludus [‡]	3.07 (1.58)	3.02 (0.98)
Love style Storge [‡]	5.49 (1.08)	5.93 (1.18)
Love style Pragma [‡]	4.68 (1.11)	4.72 (1.34)
Love style Mania [‡]	3.94 (1.48)	3.88 (1.16)
Love style Agape [‡]	6.68 (0.87)	6.84 (1.09)

Table S1. Relationship characteristics (DSC and RPL study)

*There was no significant difference in any relationship measure between the OXT and PLC sessions (all P > 0.30). [†]Love in the relationship was measured with the Passionate Love Scale (PLS).

[‡]Different love styles were assessed using a German version of Lee's love styles (MEIL).

				MN	l coordin	ates
Region	Right/left	Cluster size (voxels)	Z-score	x	у	z
OXT						
Thalamus/hypothalamus*	R	77	4.84	3	-4	-2
Posterior midbrain	L/R	67	4.29	0	-28	1
Middle occipital gyrus	R	169	4.10	33	-85	16
Anterior cingulate cortex	L/R	147	4.09	0	35	10
Cuneus	L	44	3.99	-21	-94	4
Medial frontal gyrus	R	32	3.87	12	53	7
Insula	L	27	3.67	-30	17	-14
Inferior frontal gyrus (tri)	L	25	3.74	-33	32	4
Calcarine	R	11	3.42	18	-91	-2
PLC						
Precuneus	L	56	4.55	-18	-61	40
Middle occipital gyrus	L	235	4.43	-39	-79	-2
Middle occipital gyrus	R	62	4.18	21	-94	4
Inferior frontal gyrus (tri)	R	50	4.10	45	35	10
Precuneus	R	51	4.07	27	-55	49
Middle temporal gyrus	R	23	3.39	36	-73	28

Table S2. Areas showing significantly greater activation for the partner compared with unfamiliar controls (DSC study)

The whole-brain analysis was thresholded at an uncorrected P < 0.001 with a cluster extent threshold of k = 7 voxels.

*Significant at P < 0.05 familywise error corrected. tri, pars triangularis.

				MNI	coordina	ates
Region	Right/left	Cluster size (voxels)	Z-score	x	у	z
OXT						
Middle temporal gyrus*	L	282	5.31	-45	-58	-5
Inferior frontal gyrus	R	12	4.14	27	35	-8
Middle occipital gyrus	R	10	3.90	33	-73	1
Cuneus	L	33	3.79	-21	-88	1
Inferior temporal gyrus	R	63	3.73	42	-64	-8
Anterior cingulate cortex	L	37	3.66	-3	47	-2
Anterior cingulate cortex	L/R	14	3.48	0	35	4
Middle occipital gyrus	R	15	3.39	39	-73	22
Inferior frontal gyrus	R	8	3.30	51	38	13
PLC						
Anterior cingulate cortex	L	19	3.96	-9	53	1

Table S3. Areas showing significantly greater activation for the partner compared with unfamiliar controls (RPL study)

The whole-brain analysis was thresholded at an uncorrected P < 0.001 with a cluster extent threshold of k = 7 voxels.

*Significant at P < 0.05 familywise error corrected.

PNAS PNAS

Table S4. State measurement of anxiety, mood, and attention (DSC study)

Variable	OXT group ($n = 20$) [mean (±SD)]	PLC group ($n = 20$) [mean (±SD)]	t	Р
STAI–pre	32.79 (4.66)	33.26 (6.85)	-0.36	0.71
STAI–post	34.21 (5.89)	34.74 (6.81)	-0.60	0.56
PANAS-positive-pre	29.63 (5.55)	28.32 (5.64)	1.27	0.22
PANAS-positive-post	26.53 (7.03)	25.21 (6.2)	1.52	0.15
PANAS-negative-pre	11.25 (1.25)	12.45 (6.92)	-0.79	0.44
PANAS-negative-post	11.37 (1.54)	11.32 (2.14)	0.14	0.89
D2	224.53 (33.42)	235.21 (34.62)	-1.99	0.06

State anxiety before and after the experiment was assessed using the STAI. Mood before and after the experiment was assessed using the PANAS. Attention performance after the experiment was assessed using the D2 (n = 19).

	Table S5.	State measurement	of anxiety,	mood, and	attention	(RPL stuc	ly)
--	-----------	-------------------	-------------	-----------	-----------	-----------	-----

SANG SANG

Variable	OXT group ($n = 20$) [mean (±SD)]	PLC group ($n = 20$) [mean (±SD)]	t	Р
STAI–pre	31.65 (5.60)	30.55 (5.29)	1.27	0.22
STAI–post	31.47 (5.59)	31.11 (4.85)	0.39	0.70
PANAS-positive-pre	28.80 (6.00)	28.25 (5.53)	0.48	0.64
PANAS-positive-post	28.20 (5.66)	28.40 (6.55)	-0.22	0.83
PANAS-negative-pre	10.85 (1.23)	11.90 (3.54)	-1.47	0.16
PANAS-negative-post	11.20 (3.09)	10.75 (1.33)	0.96	0.35
D2	215.55 (46.10)	221.25 (47.30)	-0.80	0.43

State anxiety before and after the experiment was assessed using the STAI. Mood before and after the experiment was assessed using the PANAS. Attention performance after the experiment was assessed using the D2 (n = 19).

Variable	DSC study [mean (±SD)]	RPL study [mean (±SD)]
Age, y	25.05 (3.25)	26.55 (3.76)
Years of education	17.35 (2.28)	17.20 (2.31)
RAVLT		
Trials 1–5*	62.6 (8.24)	
Trial 6 retention [†]	13.3 (2.0)	
Trial 7 delayed recall [‡]	13.35 (1.87)	
LPS-4	30.5 (3.87)	
MWT-B	31.55 (2.56)	
TMT-A (s)	23.47 (6.18)	
TMT-B (s)	55.54 (10.76)	
Digit-span, forward	8.15 (1.39)	
Digit-span, backward	7.9 (2.13)	
RTI		
Simple reaction time (ms)		286.11 (24.56)
Simple movement time (ms)		341.58 (57.54)
Five-choice reaction time (ms)		313.85 (34.25)
Five-choice movement time (ms)		351.30 (60.56)
PAL		
Total errors		8.56 (6.74)
Mean errors to success		2.47 (2.45)
SWM-8		
Between errors		11.89 (13.56)
Strategy score		15.16 (4.21)
Trait anxiety [§]	31.8 (9.68)	30.85 (7.74)
BDI	3.25 (4.39)	1.90 (3.40)

Table S6. Demographics and neuropsychological performance (DSC and RPL study)

In the DSC study, verbal declarative memory performance was assessed using a German adaption of the RAVLT and included *learning performance across five trials (maximum possible score 75), [†]susceptibility to interference (maximum possible score 15), and [‡]delayed recall (maximum possible score 15). Nonverbal reasoning IQ was assessed by the LPS subtest 4 (maximum possible score 40). Verbal IQ based on lexical decisions was assessed by the MWT-B (maximum possible score 37), visual attention and task-switching was assessed using the TMT-A and TMT-B (results displayed in seconds), and working memory performance was assessed using the digit-span forward and backward test (maximum possible score 14). In the RPL study, subjects' speed of response to a visual target, visual memory, and the ability to retain spatial information were measured with the RTI, the PAL, and the SWM, respectively. Anxiety symptoms were assessed by the [§]State Trait Anxiety Inventory and depressive symptoms by the self-report BDI. BDI, Beck's Depression Scale, Version II.