

Monkeypox Virus Real Time PCR Kit



FEATURES



Approved
CE marked



Feasible
compatible to multiple
sample types



Efficient
60 min for
96 samples



Reliable
5 copies/reaction
with internal control



Fulfilling
Central African clade &
West African clade detectable

Monkeypox is a rare disease that is caused by infection with monkeypox virus. Monkeypox virus belongs to the Orthopoxvirus genus in the family Poxviridae. The Orthopoxvirus genus also includes variola virus (which causes smallpox), vaccinia virus (used in the smallpox vaccine), and cowpox virus. Monkeypox was first discovered in 1958 when two outbreaks of a pox-like disease occurred in colonies of monkeys kept for research, hence the name 'monkeypox.' In humans, the symptoms of monkeypox are similar to but milder than the symptoms of smallpox. Monkeypox begins with fever, headache, muscle aches, and exhaustion. The main difference between symptoms of smallpox and monkeypox is that monkeypox causes lymph nodes to swell (lymphadenopathy) while smallpox does not. Transmission of monkeypox virus occurs when a person comes into contact with the virus from an animal, human, or materials contaminated with the virus. The virus enters the body through broken skin (even if not visible), respiratory tract, or the mucous membranes (eyes, nose, or mouth). Laboratory tests that are used to diagnose monkeypox virus include detection of immunohistochemical testing, electron microscopy, real time polymerase chain reaction (RT-PCR), and virus isolation.

The Bioperfectus Monkeypox Virus Real Time PCR Kit is based on real-time PCR technology. Specific primers and probes are designed based on F3L gene specific areas of Monkeypox virus. In addition, the kit also contains a housekeeping gene (RNase P) as an internal control (IC) for specimen sampling and nucleic acid extraction.

• SAMPLE TYPE AND PRE-PROCESSING •

• Specimen type

Human tonsillar swab,
nasopharyngeal swab, serum,
whole blood, lesion exudate and scab.

Veterinary Use version also available!
Contact our sales rep or via
info@bioperfectus.com for inquiry.

• Pre-processing for scab or crust samples

- > Mix 0.2g-1.0g of tissue samples with sterilized quartz-sand by ratio of 1:1 (v/v) and grind thoroughly
- > Add PBS (0.01mol/L, pH 7.6-7.8) with ratio of 1:5 (w/v)
- > Freeze-thaw twice at -20°C then centrifuge at 8,000 rpm for 5 minutes at 4°C
- > Transfer 200 µL of supernatant for nucleic acid extraction

· TOTAL PCR SOLUTION ·

Nucleic Acid Extraction Reagent

Nucleic Acid Extraction Rapid Kit
(Magnetic Bead Method) SDKF60101

Viral Nucleic Acid Extraction Kit
(Magnetic Bead Method) SDK60104

Whole Blood DNA Extraction Kit
(Magnetic Beads Method) SDK60110

Nucleic Acid Extraction Kit
(Silica-Based Spin Column) SDK60102

Nucleic Acid Extraction System

SSNP-2000B, 3000A, 9600A
(32, 64, 96 channels)

SMPE-960
(3*32 channels)

SAW-96, SAW-48*
(96, 48 channels)

*coming soon

PCR instrument

Bioperfectus STC-96A,
STC-96A PLUS

Applied Biosystems 7500
QuantStudio 5

Roche LightCycler480

Bio-Rad CFX96

QIAGEN Rotor-Gene Q

Analytik Jena qTOWER

· PACKAGING ·

CATALOG

YJC70115NW-25T
YJC70115NW-50T

PRODUCT

Monkeypox Virus Real Time PCR Kit

PACKAGE

25T/Kit
50T/Kit

· KIT COMPONENTS ·

COMPONENTS

VIALS/KIT

VOLUME/25T

VOLUME/50T

PCR Reaction Mix

1

313 μ L

625 μ L

Detection Mix

1

188 μ L

375 μ L

Positive Control

1

25 μ L

50 μ L

Blank Control

1


250 μ L

250 μ L

· INSTRUCTION FOR USE ·



1
Collect Sample




3
Reaction Mix Preparation



5
Get the Result



2
Nucleic Acid Extraction



4
Run the Real-Time PCR Program

JIANGSU BIOPERFECTUS TECHNOLOGIES CO., LTD.

2nd floor, Block 10, No.188 Xinjun Ring Road, Pujiang High-Tech Park, Caohejing Development Area, Shanghai, China.

3rd & 4th floors of Building A(G19), 4th floor of Building F(G14), Ground floor of Building G20, Shuaiyu Village, Fuye village, Sixiang town, Taizhou National Medical Hi-tech Development Zone, 225300 Taizhou, Jiangsu P.R. China.

info@bioperfectus.com

www.bioperfectus.com

AUTHORIZED DISTRIBUTOR